



King's Research Portal

DOI:

[10.1016/j.toxicon.2017.07.012](https://doi.org/10.1016/j.toxicon.2017.07.012)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Jaimes-Becerra, A., Chung, R., Morandini, A. C., Weston, A., Padilla, G., Gacesa, R., Ward, M., Long, P. F., & Marques, A. C. (2017). Comparative proteomics reveals recruitment patterns of some protein families in the venoms of Cnidaria. *Toxicon*, 137, 19-26. <https://doi.org/10.1016/j.toxicon.2017.07.012>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal


Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Comparative proteomics reveals recruitment patterns of some protein families in the venoms of Cnidaria

Adrian Jaimes-Becerra, Ray Chung, André C. Morandini, Andrew J. Weston, Gabriel Padilla, Ranko Gacesa, Malcolm Ward, Paul F. Long, Antonio C. Marques



PII: S0041-0101(17)30217-9

DOI: [10.1016/j.toxicon.2017.07.012](https://doi.org/10.1016/j.toxicon.2017.07.012)

Reference: TOXCON 5671

To appear in: *Toxicon*

Received Date: 17 April 2017

Revised Date: 7 July 2017

Accepted Date: 10 July 2017

Please cite this article as: Jaimes-Becerra, A., Chung, R., Morandini, André.C., Weston, A.J., Padilla, G., Gacesa, R., Ward, M., Long, P.F., Marques, A.C., Comparative proteomics reveals recruitment patterns of some protein families in the venoms of Cnidaria, *Toxicon* (2017), doi: 10.1016/j.toxicon.2017.07.012.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Comparative proteomics reveals recruitment patterns of some protein families in
the venoms of Cnidaria**

Adrian Jaimes-Becerra^{1*}, Ray Chung²⁺, André C. Morandini¹, Andrew J. Weston³,
Gabriel Padilla⁴, Ranko Gacesa⁵, Malcolm Ward²⁺, Paul F. Long⁵⁻⁷ and Antonio C.
Marques^{1,8}

¹Departamento de Zoologia, Instituto de Biociências, Universidade de São Paulo, Rua
Matão, Trav. 14, 101, 05508-090 São Paulo, SP, Brazil.

²Proteomics Facility, Institute of Psychiatry, Psychology & Neuroscience, King's
College London, 16 De Crespigny Park, London SE5 8AF, United Kingdom.

³Mass Spectrometry Laboratory, UCL School of Pharmacy, 29/39 Brunswick Square,
London WC1N 1AX, United Kingdom.

⁴Instituto de Ciências Biomédicas, Universidade de São Paulo, Av. Professor Lineu
Prestes 1374, 05508-000 São Paulo, SP, Brazil.

⁵Faculty of Life Sciences & Medicine, King's College London, 150 Stamford Street,
London SE1 9NH, United Kingdom.

⁶Brazil Institute, King's College London, Strand, London WC2R 2LS, United Kingdom.

⁷Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, Av. Prof. Lineu
Prestes, 580, B16, 05508-000 São Paulo, SP, Brazil.

⁸Centro de Biologia Marinha, Universidade de São Paulo, Rodovia Manoel Hypólito do
Rego, km. 131,5, 11600-000 São Sebastião, Brazil.

*Corresponding author: Adrian Jaimes-Becerra, email: adrianjaimesb@usp.br

23

24 [†]Present addresses: Mr Ray Chung - Social, Genetic and Developmental Psychiatry
25 Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London,
26 De Crespigny Park, London, SE5 8AF, United Kingdom. Dr Malcolm Ward - Aulesa
27 Biosciences Limited, 44, Elgar Drive, Shefford, Bedfordshire, SG17 5RZ, United
28 Kingdom.

29

Abstract

Cnidarians are probably the oldest group of animals to be venomous, yet our current picture of cnidarian venom evolution is highly imbalanced due to limited taxon sampling. High-throughput tandem mass spectrometry was used to determine venom composition of the scyphozoan *Chrysaora lactea* and two cubozoans *Tamoya haplonema* and *Chiropsalmus quadrumanus*. Protein recruitment patterns were then compared against 5 other cnidarian venom proteomes taken from the literature. A total of 28 putative toxin protein families were identified, many for the first time in Cnidaria. Character mapping analysis revealed that 17 toxin protein families with predominantly cytolytic biological activities were likely recruited into the cnidarian venom proteome before the lineage split between Anthozoa and Medusozoa. Thereafter, venoms of Medusozoa and Anthozoa differed during subsequent divergence of cnidarian classes. Recruitment and loss of toxin protein families did not correlate with accepted phylogenetic patterns of Cnidaria. Selective pressures that drive toxin diversification independent of taxonomic positioning have yet to be identified in Cnidaria and now warrant experimental consideration.

Keywords: evolution; venom; Cnidaria; nematocysts; proteomics.

Introduction

Cnidaria is believed to be the most basal of the extant Metazoa to be venomous, having evolved since Neoproterozoic times, ~650 million years ago, long before the Cambrian radiation (Van Iten *et al.*, 2014). Cnidaria is a diverse phylum comprising over 13,500 free living or parasitic marine, freshwater and terrestrial species (Daly *et al.*, 2007 plus myxozoans by Okamura *et al.*, 2015a). Cnidaria has two major subphyla: Anthozoa and Medusozoa. Anthozoa include sea anemones and both hard and soft corals (Bridge *et al.*, 1992; Marques & Collins, 2004). Medusozoa comprise the classes Staurozoa (e.g. stalked jellyfish), Cubozoa (e.g. box jellyfish), Scyphozoa (e.g. 'true' jellyfish) and Hydrozoa (e.g. *Hydra* and relatives including several species of smaller jellyfish) (Marques & Collins, 2004; Collins *et al.*, 2006; Van Iten *et al.*, 2014). Recent molecular phylogenetic analyses have corroborated the cnidarian nature of Myxozoa, with strong support as a sister-group to Medusozoa (reviewed in Okamura *et al.* 2015b).

The most evident synapomorphy of Cnidaria is the presence of cnidae, organelles produced by the Golgi apparatus of specialised cells called cnidoblasts (Marques & Collins, 2004; Fautin, 2009; Beckmann & Özbek, 2012). Cnidae are found in various parts of the body of a cnidarian and are classified into three morphological types: nematocysts, spirocysts and ptychocysts (Östman, 2000; Özbek *et al.*, 2009). The nematocysts discharge venom and are found in all cnidarians, but are morphologically and functionally heterogeneous (David *et al.*, 2008; Fautin, 2009). In addition to prey capture and defence against predation, the venom of nematocysts may also mediate spatial intraspecific and interspecific competition (Bigger, 1980; Kass-Simon & Scappaticci, 2002).

There has been resurgence in interest surrounding the nature and evolutionary origins of cnidarian venom toxins, since the first application of high throughput tandem

mass spectrometry realised high sequence homology between cnidarian toxins and those of other venomous animals (Weston *et al.*, 2012, 2013). Many studies using genomic, transcriptomic or proteomic approaches have also realised these astonishing similarities (Balasubramanian *et al.*, 2012; Brinkman *et al.*, 2012, 2015; Li *et al.*, 2012, 2014, 2016; Gacesa *et al.*, 2015; Jouiaei *et al.*, 2015a; Macrander *et al.*, 2015, 2016; Lewis *et al.*, 2016; Ponce *et al.*, 2016, Huang *et al.*, 2016), leading to the recognition that understanding the mechanisms underpinning toxin diversification in Cnidaria could provide a platform from which the evolution of this trait in higher animals might be more fully explored (Starcevic & Long, 2013; Starcevic *et al.*, 2015; Jouiaei *et al.*, 2015b). For this to be achieved, a comprehensive inventory of toxins must first be undertaken and then mapped against different taxonomic levels from established cnidarian phylogeny. To date, studies attempting to infer evolutionary aspects of toxin recruitment in Cnidaria have suffered limited taxon sampling, but when taken together these studies have demonstrated a degree of functional recruitment of certain toxin protein families at different taxonomic levels (Rachamim *et al.*, 2014; Brinkman *et al.*, 2015; Jouiaei *et al.*, 2015b). Here, the number of venom proteomes is expanded and used with data from the literature for character mapping analysis, to establish a more complete venom assembly hypothesis between the major taxonomic classes of Cnidaria.

Material & Methods

Nematocyst proteomics: The scyphozoan *Chrysaora lactea* and two cubozoans *Tamoya haplonema* and *Chiropsalmus quadrumanus* (Figure 1) were collected with permission (SISBIO license 15031-2) on May 7th 2012 by bottom shrimp trawls (2 cm mesh size) dragged at 10 m depth along Enseada beach (Guarujá County, São Paulo State, ca. 23°43'20"S 43°23'40W). Animals were identified based on morphological characters (Morandini *et al.*, 2005; Morandini & Marques, 2010; Collins *et al.*, 2011) and intact nematocysts were isolated from excised tentacles as previously described (Weston *et al.*, 2013). To extract solubilised proteins, 1 mL of protein extraction buffer (50 mM TEAB, 0.04 % (w/v) SDS, Roche protease and phosphatase inhibitor cocktail) was added to freeze dried nematocysts. The reconstituted material was disrupted in a sonic bath (VWR, Lutterworth, UK) for 15 mins. The debris was removed by centrifugation (10,000 x g for 10 mins at 4 °C). The supernatant was decanted and the soluble protein concentration determined by Bradford assay. A volume equivalent to 15 µg of protein was made up to 15 µL in extraction buffer and added to 15 µL 2 x Laemmli sample buffer, heated for 10 mins at 95 °C and loaded onto a 4-12 % (w/v) NuPAGE gel (Life Technologies) and separated by 1D SDS-PAGE. Electrophoresis was performed in MES buffer (Life technologies) at 150 V for approximately 100 mins. The entire gel lane was then divided into 15 equal sections, excised and cut into 2 mm pieces. In-gel reduction, alkylation, and proteolytic digestion with trypsin were performed as follows: Cysteine residues were reduced with 10 mM dithiothreitol and alkylated with 55 mM iodoacetamide in 100 mM ammonium bicarbonate to form stable carbamidomethyl derivatives. Trypsin digestion was carried out overnight at 37 °C in 50 mM ammonium bicarbonate buffer and the supernatant was retained. Peptides were extracted from the gel pieces by two washes with 50 mM ammonium bicarbonate and

acetonitrile. Each wash involved shaking the gel pieces for 10 mins. The extracts were pooled with the initial digestion supernatant and then lyophilised. Lyophilised extract was reconstituted in 30 μ L of 50 mM ammonium bicarbonate buffer for LC-MS/MS.

Data analysis: Data analysis was performed as previously described (Weston *et al.*, 2013; Gacesa *et al.*, 2015) but with minor modifications. Briefly, a one search matching strategy of rawfile MS/MS data against the Tox-Prot UniProtKB/Swiss-Prot database (Jungo *et al.*, 2012) using the MASCOT search engine was first executed (Perkins *et al.*, 1999). Methionine oxidation, phosphorylation on S/T/Y, deamidation on N/D and carbamidomethyl cysteine were selected as fixed modifications. Digestion with trypsin allowed up to three missed cleavages. The data were searched with a parent ion tolerance of 5 ppm and a fragment ion tolerance of 0.5 Da. The MASCOT result files were next uploaded into Scaffold v4.3.4 (Proteome Software, Portland, Oregon, USA) (Searle, 2010) and spectra corresponding to likely venom toxin peptides were manually validated for unbroken series of overlapping b-type and y-type sequence specific fragments ions, where losses consistent with the sequence were acceptable. Validated spectra (Figures S1-S3) corresponding to peptides with predicted venom toxin functions were next distinguished from peptides with likely other non-toxic physiological functions using 'ToxClassifier' (Gacesa *et al.*, 2016). This is a suite of machine learning based classifiers that provide consistent discrimination of toxins from non-toxin peptide sequences with > 99 % accuracy by performing BLAST and HMM comparisons against the Tox-Prot UniProtKB/Swiss-Prot (Jungo *et al.*, 2012), UniProt Trembl (The UniProt consortium, 2017) and NR (NCBI Resource Coordinators, 2016) databases.

Character mapping analysis: In addition to the data acquired in this study, putative toxins from other cnidarians described in the literature were also included to enhance

the dataset. These putative toxins were from the anthozoans *Anemonia viridis* (Rachamim *et al.*, 2014) and *Acropora digitifera* (Gacesa *et al.*, 2015), the hydrozoans *Olindias sambaquiensis* (Weston *et al.*, 2013) and *Hydra magnipapillata* (Rachamim *et al.*, 2014) and, the scyphozoan *Aurelia aurita* (Rachamim *et al.*, 2014). The putative toxins from the combined data set were assigned to venom toxin protein families using established KEGG ontology. Data were coded in a matrix as presence (1) or absence (0) of each toxin protein family in each species. Reconstruction of ancestral states at different nodes on an accepted taxonomic tree of Cnidaria (Marques & Collins, 2004; Collins *et al.*, 2006) was performed using Mesquite version 3.04 (Maddison & Maddison, 2015) with the parsimony criterion for the model unordered. In addition, the matrix of presences and absences of toxin protein families was used to infer a phylogenetic pattern based on the parsimony criterion.

Results

Comparative proteomics of toxin protein families: The putative toxin proteomes of nematocysts for the 3 species experimentally acquired in this study are given in Table 1. The toxin protein families from 5 species taken from the literature are given in Table S1. A total of 28 toxin protein families were identified from the nematocyst proteomes of the 8 species studied and are shown in Figure 2. Nine (~33 %) out of the 28 toxin protein families were shared by all the four classes of cnidarians. These 9 protein families were conotoxins O, CRISP, latrotoxin, lipase, metalloproteinase, phospholipases A₂ (PLA₂), phospholipases D, CS $\alpha\beta$ potassium channel blocker, and CS $\alpha\beta$ sodium channel inhibitor. Nineteen protein toxin families were not distributed across all classes (Figure 2). These included three families of pore forming toxins, which were the jellyfish toxin family-like proteins (JFTs) found to be restricted to the sister classes

Cubozoa and Scyphozoa; the actinoporins found in the classes Anthozoa, Hydrozoa and Cubozoa, and laticins found in the classes Anthozoa and Scyphozoa. The ficolins and snaclec belong to the lectin families of toxins and were limited to the Scyphozoa Anthozoa, and Hydrozoa. Peptides with similarity to three families of neurotoxins were also taxonomically restricted (Figure 2), these were the kunitz type family detected in Anthozoa and Scyphozoa, the calcium channel blocker Huwentoxin-1 reported here for the first time but solely in medusozoans, and snake three finger found in all classes except Cubozoa. Likewise, peptidase S1, flavin amino-oxidase and glycosyl hydrolase 56 families were identified in all classes except Cubozoa. Complement C3 family-like proteins were identified in the Hydrozoa and Scyphozoa. MAC-PF family-like proteins were identified in the Hydrozoa and Cubozoa. The presence of translationally controlled tumour like proteins (TCTP) was identified in the venom proteome from both Anthozoa and Hydrozoa.

Evolution of the cnidarian venom arsenal: Recruitment patterns of putative toxin protein families (Figure 3 and Table S3) were inferred using a presence and absence matrix (Table S2). This recruitment pattern indicated that venom of Medusozoa and Anthozoa ancestors might have been composed of at least seventeen types of protein toxin families (Figure 3 and Figure S4i). After separation of the ancestral lineage into Anthozoa and Medusozoa, some putative toxins were lost (or not expressed) in some clades. For example, the TCTP family was not present in the Cubozoa and Scyphozoa. Similarly, the actinoporin toxin protein family was lost from Scyphozoa. Unlike the other classes, the species of Cubozoa examined demonstrated large losses. Nine toxin protein families might have been recruited by a single clade after the split Anthozoa-Medusozoa (Figure 3 and Figure S4ii). Three families of cytolytic toxins (MAC-PF, ficolin lectin and JFTs) appear to have been recruited into Medusozoa after the basal diversification event into the venom of Hydrozoa, Scyphozoa, and Cubozoa (Figure 3). Equally, two families of neurotoxins ShK-like potassium channel and sea anemone sodium channel modulator appear to have been recruited into Anthozoa only. The laticin and kunitz-type toxin protein families, might have been recruited independently (i.e., by convergence) into the venom of Anthozoa/Scyphozoa (Figure 3 and Figure S4iii). Phylogenetic analysis of the presence and absence matrix gave a topology of (Cubozoa(Anthozoa(Hydrozoa,Scyphozoa))), which disagreed with the more accepted phylogeny of Cnidaria (Anthozoa(Hydrozoa(Cubozoa,Scyphozoa))).

Discussion

The putative toxin component of nematocyst proteomes for 3 out of the 8 species examined (*Chrysaora lactea*, *Tamoya haplonema*, and *Chiropsalmus quadrumanus*) are described in this study for the first time (Table 1). Venom data from 3 other species

(*Anemonia viridis*, *Hydra magnipapillata*, and *Aurelia aurita*) were published elsewhere (Rachamim *et al.*, 2014) and reassessed in this study. These data were combined with our own previously published putative nematocyst toxin proteomes from *Acropora digitifera* (Gacesa *et al.*, 2015) and *Olindias sambaquiensis* (Weston *et al.*, 2013). Altogether, the data from this study has supported previous research that Anthozoa and Medusozoa have complex venom composition comprising multiple protein families (Rachamim *et al.*, 2014; Jouiaei *et al.*, 2015c) (Figures 2 and 3). We highlight that, although transcriptomes and proteomes from other species of cnidarians have also been published (Moran *et al.*, 2013; Jouiaei *et al.*, 2015a, Ponce *et al.*, 2016, Macrander *et al.*, 2016), our analysis focused on those species for which we had access to raw proteomics MS/MS data which could be analysed using identical bioinformatics methods, ensuring results were fully comparable. Our study was conservative, being restricted to putative toxin annotation in the expressed proteome and did not include a study of transcriptomes. This was because not all the transcripts that contributed to transcriptome diversity would equally be likely to be translated (if at all) and have contributed to protein diversity. Hence, correlation between sequences annotated as putative toxins in the transcriptome and proteome would not have been straightforward given the difficulty in differentiating sequences with toxic and other physiological functions. Future work to overcome this impediment will require the acquisition of genome sequence onto which other sequence data can be mapped (Gacesa *et al.*, 2015).

Comparative venom proteomic analysis from different Cnidaria classes

Our comparative proteomics data of putative venom toxins indicated that nearly half of the protein toxin families were distributed across all of the cnidarian classes studied (Figure 2). The biological activities of some of these toxin families are of note,

for example, PLA₂ toxins have thus far only been identified with haemolytic activity in cnidarians (Hessinger & Lenhoff, 1976; Grotendorst & Hessinger, 2000; Anderluh & Maček, 2002; Talvinen & Nevalainen, 2002; Nevalainen *et al.*, 2004; Razpotnik *et al.*, 2010), although neurotoxic and myotoxic activities as well as non-toxic physiological functions have also been widely reported in other venomous animals (Fry *et al.*, 2009; Six & Dennis, 2000). Likewise, phospholipase D family proteins isolated from cnidarian venoms have been reported to exhibit necrotic activity (Burke, 2002; Uri *et al.*, 2005), with homologs also recently identified in the giant jellyfish *Cyanea capillata* (Liu *et al.*, 2015). Most of the metalloproteinases identified in this study belonged to the zinc metalloproteinase family. This family of toxins is an important component found in the venoms of many terrestrial animals such as centipedes, snakes and ticks (Fry *et al.*, 2009; Undheim *et al.*, 2014), with diverse biological activities culminating in haemorrhage and tissue necrosis in the target prey following envenomation (Fox & Serrano, 2005; da Silveira *et al.*, 2007). Transcriptomic and proteomic studies have identified zinc metalloprotease in venoms of the scyphozoans *Stomolophus meleagris*, *Cyanea capillata*, and *Cyanea nozakii* (Li *et al.*, 2014, 2016; Liu *et al.*, 2015), the cubozoan *Chironex fleckeri* (Brinkman *et al.*, 2015; Jouiaei *et al.*, 2015a) and the anthozoan *Anthopleura elegantissima* (Macrander *et al.*, 2015). A study of metalloproteases from the scyphozoan *Nemopilema nomurai*, *Rhopilema esculenta*, *Cyanea nozakii*, and *Aurelia aurita* confirmed the necrotic toxicity of these enzymes (Lee *et al.*, 2011). Both sodium and potassium ion channel inhibitors were identified in representatives of all of the classes examined. These two types of neurotoxins have been widely studied in Anthozoa, especially sea anemones and comprise the largest number of toxins so far recorded in public databases for Cnidaria (Moran *et al.*, 2009; Šuput, 2009; Turk & Kem, 2009; Frazão *et al.*, 2012; Jouiaei *et al.*, 2015c; Macrander *et al.*,

2015; Mariottini *et al.*, 2015). Neurotoxic effects have been identified in scyphozoans such as *Cyanea nozakii* (Feng *et al.*, 2010), *Cyanea capillata* (Helmholz *et al.*, 2012), and *Pelagia noctiluca* (Pang *et al.*, 1993; Morabito *et al.*, 2012) and, in cubozoans such as *Carukia barnesi* (Winkel *et al.*, 2005) and *Malo kingi* (Gershwin, 2007). In this study, we identified two putative types of sodium and potassium ion channel inhibitors (Figure 2, Table S2). ShK-like potassium channel and sea anemone sodium channel modulator were only found in a single Anthozoan species, *Anemonia viridis* (a sea anemone). It should be noted that this was the only species of sea anemone analysed in this study. Both sodium and potassium putative ion channel inhibitors have been found exclusively in sea anemones (Moran *et al.*, 2009; Diochot and Lazdunski, 2009). CS $\alpha\beta$ potassium channel blocker and CS $\alpha\beta$ sodium channel inhibitor were found in all of the species of cnidarians analysed including another anthozoan, *Acropora digitifera* and have sequence homology to sodium and potassium channel blockers of scorpions. This observation might highlight a rare example of mechanistic convergence whereby sodium and potassium ion channel inhibitors appeared on two separate occasions within the cnidarians. Convergent evolution of these toxins in scorpions and sea anemones has been previously reported and although these toxin protein families are structurally different, functional mapping studies have shown similarities in the binding cores (Gasparini *et al.*, 2004). Further species sampling is required to substitute these observations which are based here on a single MS/MS event in the anthozoa *Acropora digitifera*, the hydrozoa *Hydra magnipapillata*, the scyphozoan *Aurelia aurita* and the cubozoans *Chiropsalmus quadrumanus* and *Tamoya haplonema*. It should also be noted that the names given to each putative ion channel inhibitors were used to distinguish between the two different possible origins of the putative sodium and potassium ion channel inhibitors identified in this study. Another family of neurotoxins were the

CRISP type toxins, which again were found in all classes of cnidarians studied herein. This toxin protein family has widely been reported in cnidarian venoms (Brinkman *et al.*, 2015; Ponce *et al.*, 2016; Lewis *et al.*, 2016) and attributed many biological functions.

Just over half of the toxins protein families identified in this study were restricted to certain cnidarian classes only (Figure 2). Hyaluronidase-like proteins were found in all classes of cnidarians except Cubozoa, but these proteins are common and have non-toxic physiological function in many non-venomous animals. It is feasible that such proteins are likely recruited into venoms not as toxins, but as adjuvants to increase tissue permeability (Kemparaju & Girish, 2006; Fry *et al.*, 2009). Non-toxic peptides and proteins present in nematocysts that may function in toxin maturation, toxin trafficking and delivery, or as self-defence mechanisms against the biological activities of the venom have received little study and perhaps warrant closer inspection. Likewise, the peptidase S1 family was also detected in all cnidarian classes studied except Cubozoa. This family is part of the group of serine protease inhibitors that is widely distributed in other marine venomous animals including marine cone snails and cephalopods (Mourão & Schwartz, 2013), as well as terrestrial reptiles (Fry *et al.*, 2009). Recently, serine protease homologs were identified in the transcriptome of the sea anemone *Anthopleura elegantissima* (Macrander *et al.*, 2015). However, the biological activity of the S1 peptidase family of toxins has yet to be confirmed in cnidarians. It is unclear why proteins commonly associated with innate immune responses are also apparently widely distributed in cnidarian venoms. For example, MAC-PF-like toxins have also been identified in sea anemones (Nagai *et al.*, 2002b; Oshiro *et al.*, 2004) and were recently annotated in the transcriptomes and proteomes of Hydrozoa and Scyphozoa (Rachamim *et al.*, 2014). Likewise, the actinoporins are pore-

forming toxins were found in the proteomes of Anthozoa and Cubozoa classes. These cytolysins have also been identified in transcriptome sequences and biological activity recorded in nematocyst venom extracts of various *Hydra* species (Hydrozoa) (Hwang *et al.*, 2007; Glasser *et al.*, 2014).

Assembly of the cnidarian venom proteome

To date, only one previous study has been published that used similar approaches to those described here to investigate evolutionary aspects of toxin recruitment in Cnidaria (Rachamim *et al.*, 2014). In this previous study, the kunitz type family of toxins was only found in Anthozoa. In the study present here, this family of toxins was found in Anthozoa, but was also identified in Scyphozoa. In the study of Rachamim *et al.*, (2014), the PLA2 family of toxin proteins were only found in species of Scyphozoa and Hydrozoa. In our analyses, PLA2 were present in all the Cnidaria classes studied and hence, most likely arose as a recruitment event at the base ancestor of the Cnidaria. It should be noted that PLA2 like proteins have also been identified in recent studies of Anthozoan venoms (Macrander *et al.*, 2015, 2016). Differences in the recruitment patterns between studies might be explained because of the low number of species sampled. The extent of comparison groups (all Cnidaria) in light of the sparseness of data at terminals in the analysis is a concern, for example, no data is currently available on the toxin complement of venoms from the Staurozoa or Myxozoa (Marques & Collins, 2004; Okamura *et al.*, 2015a). Based on the data presented, many of the neurotoxic and cytotoxic protein toxin families might have been recruited into the venom proteome early in cnidarian evolution, before the first major radiation in this phylum around 800 million years ago (Park *et al.*, 2012; Van Iten *et al.*, 2017).

The approaches used in this study were very conservative, with analyses based exclusively on putative toxin protein families found in each proteomic profile and not specific toxin peptides or proteins. These proteomic profiles can be considered phenotypes, or a "morphological representation" of the venom, allowing variation in the toxin complement to be evaluated. For example, in this study JFTs were found only in the venoms of Scyphozoa and Cubozoa. However, previous studies have demonstrated JFTs encoded in the genome and expressed in the proteome of Hydrozoa, and in the transcriptome of the anthozoan *Anemonia viridis* (Rachamim *et al.*, 2014). Few reports in the literature have documented variation in toxin composition of venom at taxonomic level in the phylum Cnidaria (Orts *et al.*, 2013; Rachamim *et al.*, 2014), and certainly there have been no studies that have attempted to put into context what the biological consequences of venom variation might be (Gacesa *et al.*, 2015; Knittell *et al.*, 2016). The difference between the two phylogenetic patterns (accepted vs inferred using the presence and absence matrix) found in this study could be due to various ecological factors that need to be investigated in future studies. But increased sampling and analysis at different taxonomic levels is a priority in order to identify the influence of history and ecology in the origin of these contrasting patterns.

Conclusions

Venom composition of Medusozoa and Anthozoa are different, with cytolytic toxin protein families slightly more abundant in Medusozoa. When only toxin protein family composition was used for phylogenetic inference, the resulting topology (Cubozoa(Anthozoa(Hydrozoa,Scyphozoa))) did not match the classic published phylogeny (Anthozoa(Hydrozoa(Cubozoa,Scyphozoa))). Understanding the functional context (environment versus morphological form) that may drive expression of toxins in

Cnidaria requires future experimental consideration, including wider taxonomic sampling.

Acknowledgments

We are indebted to Dr David Morganstern, Prof Tamar Lotan and Prof Daniel Sher for making available their mass spectrometry data. We extend our thanks to the Santos Family for providing logistic support of shipboard operations. We are also grateful for onshore technical assistance of the staff at CEBIMar. This is a contribution to the NP-BioMar program at the Universidade de São Paulo.

Funding

This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo [grant numbers 2010/52324-6, 2010/06927-0, 2011/50242-5, 2013/50484-4, 2010/50174-7], the Conselho Nacional de Desenvolvimento Científico e Tecnológico [grant numbers 562143/2010-6, 477156/2011-8, 305805/2013-4, 445444/2014-2, 301039/2013-5], the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior [grant number 236.507.518-52], the Universidade de São Paulo [grant number 13.1.1502.9.8] and the Medical Research Council [grant number G82144A].

References

Anderluh G, Maček, P. 2002. Cytolytic peptide and protein toxins from sea anemones (Anthozoa: Actiniaria). *Toxicon* 40:111–124 DOI:10.1016/S0041-0101(01)00191-X.

- 377 Balasubramanian PG, Beckmann A, Warnken U, Schnölzer M, Schüler A, Bornberg-
378 Bauer E, Holstein TW, Özbek S. 2012. Proteome of *Hydra* nematocyst. The Journal of
379 Biological Chemistry 287:9672–9681 DOI: 10.1074/jbc.M111.328203.
- 380 Beckmann A, Özbek S. 2012. The nematocyst: A molecular map of the cnidarian
381 stinging organelle. The International Journal of Developmental Biology 56:577–582
382 DOI: 10.1387/ijdb.113472ab.
- 383 Bigger CH. 1980. Interspecific and intraspecific acrorhagial aggressive behaviour
384 among sea anemones: a recognition of self and not self. The Biological Bulletin
385 159:117–134 DOI: 10.2307/1541013.
- 386 Bridge D, Cunningham CW, DeSalle R, Buss LW. 1992. Class-level relationships in the
387 phylum Cnidaria: Evidence from mitochondrial genome structure. Proceedings of the
388 National Academy of Sciences of the United States of America 89:8750–8753
389 DOI:10.1073/pnas.89.18.8750.
- 390 Brinkman DL, Aziz A, Loukas A, Potriquet J, Seymour J, Mulvenna J. 2012. Venom
391 proteome of the box jellyfish *Chironex fleckeri* PLoS ONE 7: e47866
392 DOI:10.1371/journal.pone.0047866.
- 393 Brinkman DL, Jia X, Potriquet J, Kumar D, Dash D, Kvaskoff D, Mulvenna J. 2015.
394 Transcriptome and venom proteome of the box jellyfish *Chironex fleckeri*. BMC
395 Genomics 16 DOI: 10.1186/s12864-015-1568-3
- 396 Burke WA. Cnidarians and human skin. 2002. Dermatologic Therapy 15:18–25 DOI:
397 10.1046/j.1529-8019.2002.01508.x.

- 398 Collins AG, Bentlage B, Gillan W, Lynn TH, Morandini AC, Marques AC. 2011.
399 Naming the Bonaire banded box jelly, *Tamoya ohboya*, n. sp. (Cnidaria: Cubozoa:
400 Carybdeida: Tamoyidae). Zootaxa 2753:53–68 DOI: [http://dx.doi.org/10.11646/%25x](http://dx.doi.org/10.11646/zootaxa.2753.1.2)
- 401 Collins AG, Schuchert P, Marques AC, Jankowski T, Medina M, Schierwater B. 2006.
402 Medusozoan phylogeny and character evolution clarified by new large and small
403 subunit rDNA data and an assessment of the utility of phylogenetic mixture models.
404 Systematic Biology 55:97–115 DOI: 10.1080/10635150500433615
- 405 Da Silveira RB, Wille ACM, Chaim OM, Appel MH, Silva DT, Franco CRC, Toma L,
406 Mangili OC, Gremski W, Dietrich CP, Nader HB, Veiga SS. 2007. Identification,
407 cloning, expression and functional characterization of an astacin-like metalloprotease
408 toxin from *Loxosceles intermedia* (Brown spider) venom. Biochemical Journal
409 406:355–363 DOI: 10.1042/BJ20070363.
- 410 Daly M, Brugler MR, Cartwright P, Collins AG, Dawson MN, Fautin DG, France SC,
411 McFadden CS, Opresko DM, Rodriguez E, Romano SL, Stake JL. 2007. The phylum
412 Cnidaria: A review of phylogenetic patterns and diversity 300 years after Linnaeus.
413 Zootaxa 1668:127–182.
- 414 David CN, Özbek S, Adamczyk P, Meier S, Pauly B, Chapman J, Hwang JS, Gojobori
415 T, Holstein TW. 2008. Evolution of complex structures: minicollagens shape the
416 cnidarian nematocyst. Trends in Genetics. 24:431–438 DOI:
417 <http://dx.doi.org/10.1016/j.tig.2008.07.001>.
- 418 Diochot S, Lazdunski M. 2009. Sea anemone toxins affecting potassium channels. In:
419 Fusetani N, Kem W, eds. Marine toxins as research tools. Cham: Springer, 99–122.
- 420 Fautin DG. 2009. Structural diversity, systematics, and evolution of cnidae. Toxicon
421 54:1054–1064 DOI:10.1016/j.toxicon.2009.02.024

- 422 Feng J, Yu H, Xing R, Liu S, Wang L, Cai S, Li P. 2010. Isolation and characterization
423 of lethal proteins in nematocyst venom of the jellyfish *Cyanea nozakii* Kishinouye.
424 *Toxicon* 55:118–125 DOI:10.1016/j.toxicon.2009.07.008
- 425 Fox JW, Serrano SMT. 2005. Structural considerations of the snake venom
426 metalloproteinases, key members of the M12 reprotolysin family of metalloproteinases.
427 *Toxicon* 45: 969–985 DOI:10.1016/j.toxicon.2005.02.012.
- 428 Frazão B, Vasconcelos V, Antunes A. Sea anemone (Cnidaria, Anthozoa, Actiniaria)
429 toxins: An overview. 2012. *Marine Drugs* 10:1812–1851 DOI:10.3390/md10081812.
- 430 Fry BG, Roelants K, Champagne DE, Scheib H, Tyndall JD, King GF, Nevalainen TJ,
431 Norman J, Lewis RJ, Norton RS, Renjifo C, de la Vega RCR. 2009. The toxicogenomic
432 multiverse: convergent recruitment of proteins into animal venoms. *Annual Reviews of*
433 *Genomics and Human Genetics* 10:483–511 DOI:
434 10.1146/annurev.genom.9.081307.164356.
- 435 Gacesa R, Chung R, Dunn SR, Weston A, Jaimes-Becerra A, Marques AC, Morandini
436 A, Hranueli D, Starcevic A, Ward M, Long PF. 2015. Gene duplications are extensive
437 and contribute significantly to the toxic proteome of nematocysts isolated from
438 *Acropora digitifera* (Cnidaria: Anthozoa: Scleractinia). *BMC Genomics* 16:774 DOI:
439 10.1186/s12864-015-1976-4.
- 440 Gacesa R, Barlow D, Long P. 2016. Machine learning can differentiate venom toxins
441 from other proteins having non-toxic physiological functions. *PeerJ Computer Science*
442 2:e90 DOI 10.7717/peerj-cs.90.
- 443 Gasparini S, Gilquin B, Ménez A. 2004. Comparison of sea anemone and scorpion
444 toxins binding to Kv1 channels: an example of convergent evolution. *Toxicon* 43:901–
445 908 DOI: 10.1016/j.toxicon.2004.03.029.

- 446 Gershwin LA. 2007. *Malo kingi*: A new species of Irukandji jellyfish (Cnidaria:
447 Cubozoa: Carybdeidae), possibly lethal to humans, from Queensland, Australia.
448 Zootaxa 68:55–68.
- 449 Glasser E, Rachamim T, Aharanovich D, Sher D. 2014. Hydra actinoporin-like toxin-1,
450 an unusual hemolysin from the nematocyst venom of *Hydra magnipapillata* which
451 belongs to an extended gene family. Toxicon 91:103–113
452 DOI:<https://doi.org/10.1016/j.toxicon.2014.04.004>
- 453 Grotendorst GR, Hessinger DA. 2000. Enzymatic characterization of the major
454 phospholipase A2 component of sea anemone (*Aiptasia pallida*) nematocyst venom.
455 Toxicon 38:931–943 DOI:10.1016/S0041-0101(99)00206-8.
- 456 Helmholz H, Wiebring A, Lassen S, Ruhnau C, Schuett C, Prange A. 2012. Cnidom
457 analysis combined with an *in vitro* evaluation of the lytic, cyto- and neurotoxic potential
458 of *Cyanea capillata* (Cnidaria: Scyphozoa). Scientia Marina 76: 339–348
459 <http://dx.doi.org/10.3989/scimar.03381.16E>.
- 460 Hessinger DA, Lenhoff HM. 1976. Mechanism of haemolysis induced by nematocyst
461 venom: Roles of phospholipase A and direct lytic factor. Archives of Biochemistry and
462 Biophysics 173:603–613 DOI:10.1016/0003-9861(76)90297-6.
- 463 Huang C. Morlighem JE. Zhou H. Lima EP. Gomes PB. Cai J. Lou I. Perez C. Lee SM.
464 Rádis-Baptista G. 2016. The Transcriptome of the zoanthid *Protopalythoa variabilis*
465 (Cnidaria, Anthozoa) predicts a basal repertoire of toxin-like and venom-auxiliary
466 polypeptides. Genome Biology and Evolution 8:3045–3064 doi:10.1093/gbe/evw204.
- 467 Hwang JS, Ohyanagi H, Hayakawa S, Osato N, Nishimiya-Fujisawa C, Ikeo K, David
468 C, Fujisawa T, Gojobori T. 2007. The evolutionary emergence of cell type-specific
469 genes inferred from the gene expression analysis of *Hydra*. Proceedings of the National

- 470 Academy of Sciences of the United States of America 104: 14735–14740 DOI:
471 10.1073/pnas.0703331104.
- 472 Jouiaei M, Casewell N, Yanagihara A, Nouwens A, Cribb B, Whitehead D, Jackson T,
473 Ali S, Wagstaff S, Koludarov I, Alewood P, Hansen J, Fry B. 2015a. Firing the sting:
474 Chemically induced discharge of cnidae reveals novel proteins and peptides from box
475 jellyfish (*Chironex fleckeri*) venom. *Toxins* 7:936–950 DOI:10.3390/toxins7030936.
- 476 Jouiaei M, Sunagar K, Gross AF, Scheib H, Alewood PF, Moran Y, Fry BG. 2015b.
477 Evolution of an ancient venom: recognition of a novel family of cnidarian toxins and
478 the common evolutionary origin of sodium and potassium neurotoxins in sea anemone.
479 *Molecular Biology and Evolution* 32:1598–1610 DOI: 10.1093/molbev/msv050.
- 480 Jouiaei M, Yanagihara AA, Madio B, Nevalainen TJ, Alewood PF, Fry BG. 2015c.
481 Ancient venom systems: A review on Cnidaria toxins. *Toxins* 7:2251–2271 DOI:
482 10.3390/toxins7062251.
- 483 Jungo F, Bougueleret L, Xenarios I, Poux S. 2012. The UniProtKB/Swiss-Prot Tox-Prot
484 program: a central hub of integrated venom protein data. *Toxicon* 60:551–557 DOI:
485 10.1016/j.toxicon.2012.03.010.
- 486 Kass-Simon G, Scappaticci J. 2002. The behavioural and developmental physiology of
487 nematocysts. *Canadian Journal Zoology* 80:1772–1794 DOI: 10.1139/z02-135.
- 488 Kemparaju K, Girish KS. 2006. Snake venom hyaluronidase: A therapeutic target. *Cell*
489 *Biochemistry and Function* 24:7–12 DOI: 10.1002/cbf.1261.
- 490 Knittel P, Long PF, Brammall L, Marques AC, Almeida MT, Padilla G, Moura-da-Silva
491 AM. 2016. Characterising the enzymatic profile of crude tentacle extracts from the
492 South Atlantic jellyfish *Olindias sambaquiensis* (Cnidaria: Hydrozoa). *Toxicon* 119:1–7
493 DOI: <http://dx.doi.org/10.1016/j.toxicon.2016.04.048>.

- 494 Lee H, Jung ES, Kang C, Yoon WD, Kim JS, Kim E. 2011. Scyphozoan jellyfish venom
 495 metalloproteinases and their role in the cytotoxicity. *Toxicon* 58:277–284
 496 DOI:10.1016/j.toxicon.2011.06.007.
- 497 Lewis-Ames C, Ryan J, Bely A, Cartwright P, Collins A. 2016. A new transcriptome and
 498 transcriptome profiling of adult and larval tissue in the box jellyfish *Alatina alata*: an
 499 emerging model for studying venom, vision and sex. *BMC Genomics* 17:650
 500 DOI:10.1186/s12864-016-2944-3.
- 501 Li R, Yu H, Xing R, Liu S, Qing Y, Li K, Li B, Meng X, Cui J, Li P. 2012. Application
 502 of nanoLC-MS/MS to the shotgun proteomic analysis of the nematocyst proteins from
 503 jellyfish *Stomolophus meleagris*. *Journal of Chromatography B* 899:86–95
 504 DOI:10.1016/j.jchromb.2012.05.006.
- 505 Li R, Yu H, Xue W, Yue Y, Liu S, Xing R, Li P. 2014. Jellyfish venomomics and venom
 506 gland transcriptomics analysis of *Stomolophus meleagris* to reveal the toxins associated
 507 with sting. *Journal of Proteomics* 106:17–29 DOI:10.1016/j.jprot.2014.04.011.
- 508 Li R, Yu H, Yue Y, Liu S, Xing R, Chen X, Li P. 2016. Combined proteomics and
 509 transcriptomics identifies sting-related toxins of Jellyfish *Cyanea nozakii*. *Journal of*
 510 *proteomics* 148:57–64 DOI:http://dx.doi.org/10.1016/j.jprot.2016.07.023.
- 511 Liu G, Zhou Y, Liu D, Wang Q, Ruan Z, He Q, Zhang L. 2015. Global transcriptome
 512 analysis of the tentacle of the jellyfish *Cyanea capillata* using deep sequencing and
 513 expressed sequence tags: Insight into the toxin- and degenerative disease-related
 514 transcripts. *PLoS ONE* 10:e0142680
 515 DOI:http://dx.doi.org/10.1371/journal.pone.0142680.

- 516 Macrander J, Brugler MR, Daly M. 2015. A RNA-seq approach to identify putative
517 toxins from acrorhagi in aggressive and non-aggressive *Anthopleura elegantissima*
518 polyps. BMC Genomics 16:221 DOI: 10.1186/s12864-015-1417-4.
- 519 Macrander J, Broe M, Daly M. 2016. Tissue-specific venom composition and
520 differential gene expression in sea anemones. Genome Biology and Evolution
521 DOI:10.1093/gbe/evw155.
- 522 Maddison WP, Maddison DR. 2015. Mesquite: a modular system for evolutionary
523 analysis. Version 3.04 <http://mesquiteproject.org>.
- 524 Mariottini GL, Bonello G, Giacco E, Pane L. 2015. Neurotoxic and neuroactive
525 compounds from Cnidaria: Five decades of research.... and more. Central Nervous
526 System Agents in Medicinal Chemistry 15:74–80 DOI:
527 10.2174/1871524915666150309141900.
- 528 Marques AC, Collins AG. 2004. Cladistic analysis of Medusozoa and cnidarian
529 evolution. Invertebrate Biology 123:23–42 DOI: 10.1111/j.1744-7410.2004.tb00139.x.
- 530 Morabito R, Condello S, Currò M, Marino A, Ientile R, La Spada G. 2012. Oxidative
531 stress induced by crude venom from the jellyfish *Pelagia noctiluca* in neuronal-like
532 differentiated SH-SY5Y cells. Toxicology in Vitro 26:694–699
533 DOI:10.1016/j.tiv.2012.03.002.
- 534 Moran Y, Gordon D, Gurevitz M. 2009. Sea anemone toxins affecting voltage-gated
535 sodium channels - molecular and evolutionary features. Toxicon 54:1089–1101
536 DOI:10.1016/j.toxicon.2009.02.028
- 537 Moran Y, Praher D, Schlesinger A, Ayalon A, Tal Y, Technau U. 2013. Analysis of
538 soluble protein contents from the nematocysts of a model sea anemone sheds light on

- 539 venom evolution. *Marine Biotechnology* 15: 329–339 DOI: 10.1007/s10126-012-9491-
540 y.
- 541 Morandini AC, Ascher D, Stampar SN, Ferreira JFV. 2005. Cubozoa e Scyphozoa
542 (Cnidaria: Medusozoa) de águas costeiras do Brasil. *Iheringia Série Zoologia* 95:281–
543 294 DOI: <http://dx.doi.org/10.1590/S0073-47212005000300008>.
- 544 Morandini AC, Marques AC. 2010. Revision of the genus *Chrysaora* Péron & Lesueur,
545 1810 (Cnidaria: Scyphozoa). *Zootaxa* 2464:1–97 DOI:<http://dx.doi.org/10.11646/%25x>.
- 546 Mourão CBF, Schwartz EF. 2013. Protease inhibitors from marine venomous animals
547 and their counterparts in terrestrial venomous animals. *Marine Drugs* 11:2069–2112
548 DOI:10.3390/md11062069.
- 549 Nagai H, Oshiro N, Takuwa-Kuroda K, Iwanaga S, Nozaki M, Nakajima T. 2002. Novel
550 proteinaceous toxins from the nematocyst venom of the Okinawan sea anemone
551 *Phyllodiscus semoni* Kwietniewski. *Biochemical and Biophysical Research*
552 *Communications* 294:760–763 DOI:10.1016/S0006-291X(02)00547-8.
- 553 NCBI Resource Coordinators. 2016. Database resources of the National Center for
554 Biotechnology Information. *Nucleic Acids Research* 44:D7–D19 DOI:
555 10.1093/nar/gkv1290.
- 556 Nevalainen TJ, Peuravuori HJ, Quinn RJ, Llewellyn LE, Benzie JH, Fenner PJ, Winkel
557 KD. 2004. Phospholipase A2 in Cnidaria. *Comparative Biochemistry and Physiology*
558 *Part B: Biochemistry and Molecular Biology* 139:731–735
559 DOI:10.1016/j.cbpc.2004.09.006.

- Okamura B, Gruhl A, Reft AJ. 2015a. Cnidarian origins of Myxozoa. In: Okamura B, Gruhl A, Bartholomew JL, eds. Myxozoan Evolution, Ecology and Development. Cham: Springer, 45–68.
- Okamura B, Gruhl A, Bartholomew J. 2015b. An introduction to myxozoan evolution, ecology and development. In: Okamura B, Gruhl A, Bartholomew JL, eds. Myxozoan Evolution, Ecology and Development. Cham: Springer, 1-20.
- Orts DJB, Peigneur S, Madio B, Cassoli JS, Montandon GG, Pimenta AMC, Bicudo JEPW, Freitas JC, Zaharenko AJ, Tytgat J. 2013. Biochemical and electrophysiological characterization of two sea anemone Type 1 potassium toxins from a geographically distant population of *Bunodosoma caissarum*. *Marine Drugs* 11:655–679 DOI:10.3390/md11030655.
- Oshiro N, Kobayashi C, Iwanaga S, Nozaki M, Namikoshi M, Spring J, Nagai H. 2004. A new membrane-attack complex/perforin (MACPF) domain lethal toxin from the nematocyst venom of the Okinawan sea anemone *ActinERIA villosa*. *Toxicon* 43:225–228 DOI:10.1016/j.toxicon.2003.11.017.
- Östman C. 2000. A guideline to nematocyst nomenclature and classification, and some notes on the systematic value of nematocysts. *Scientia Marina* 64:31–46 DOI:10.3989/scimar.2000.64s131
- Özbek S, Balasubramanian PG, Holstein TW. 2009. Cnidocyst structure and the biomechanics of discharge. *Toxicon* 54:1038–1045 DOI:10.1016/j.toxicon.2009.03.006.
- Pang KA, Schawrtz MS. 1993. Guillain-Barré syndrome following jellyfish stings (*Pelagia noctiluca*). *Journal of Neurology, Neurosurgery and Psychiatry* 56:1133–1137 DOI:10.1136/jnnp.56.10.1133.

- 583 Park E, Hwang DS, Lee SJ, Song JI, Seo TK, Won YJ. 2012. Estimation of the
584 divergence times in cnidarian evolution based on mitochondrial protein-coding genes
585 and the fossil record. *Molecular Phylogenetics and Evolution* 62:329–345
586 DOI:10.1016/j.ympev.2011.10.008.
- 587 Perkins DN, Pappin DJ, Creasy DM, Cottrell JS. 1999. Probability-based protein
588 identification by searching sequence databases using mass spectrometry data.
589 *Electrophoresis* 20:3551–3567 DOI: 10.1002/(SICI)1522-
590 2683(19991201)20:18<3551::AID-ELPS3551>3.0.CO;2-2.
- 591 Ponce D, Brinkman D, Potriquet J, Mulvenna J. 2016. Tentacle transcriptome and
592 venom proteome of the pacific sea nettle, *Chrysaora fuscescens* (Cnidaria: Scyphozoa).
593 *Toxins* 8:102 DOI:10.3390/toxins8040102.
- 594 Rachamim T, Morgenstern D, Aharonovich D, Brekhman V, Lotan T, Sher D. 2014. The
595 dynamically evolving nematocyst content of an anthozoan, a scyphozoan, and a
596 hydrozoan. *Molecular Biology and Evolution* 32:740–753 DOI:
597 10.1093/molbev/msu335.
- 598 Razpotnik A, Križaj I, Šribar J, Kordiš D, Maček P, Frangež R, Kem WR, Turk T. 2010.
599 A new phospholipase A2 isolated from the sea anemone *Urticina crassicornis* - Its
600 primary structure and phylogenetic classification. *The FEBS Journal* 277:2641–2653
601 DOI: 10.1111/j.1742-4658.2010.07674.x.
- 602 Searle BC. 2010. Scaffold: a bioinformatic tool for validating MS/MS-based proteomic
603 studies. *Proteomics* 10:1265–1269 DOI: 10.1002/pmic.200900437.
- 604 Six DA, Dennis EA. 2000. The expanding superfamily of phospholipase A2 enzymes:
605 Classification and characterization. *Biochimica et Biophysica Acta (BBA) - Molecular*
606 *and Cell Biology of Lipids* 1488:1–19 DOI:10.1016/S1388-1981(00)00105-0.

- Starcevic A, Long PF. 2013. Diversification of animal venom peptides-were jellyfish amongst the first combinatorial chemists? *ChemBioChem* 14:1407–1409 DOI: 10.1002/cbic.201300305.
- Starcevic A, Moura-da-Silva AM, Cullum J, Hranueli D, Long PF. 2015. Combinations of long peptide sequence blocks can be used to describe toxin diversification in venomous animals. *Toxicon* 95:84–92 DOI:10.1016/j.toxicon.2015.01.005.
- Šuput D. 2009. *In vivo* effects of cnidarian toxins and venoms. *Toxicon* 54:1190–1200 DOI:10.1016/j.toxicon.2009.03.001.
- Talvinen KA, Nevalainen TJ. 2002. Cloning of a novel phospholipase A2 from the cnidarian *Adamsia carciniopados*. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*. 132:571–578 DOI:10.1016/S1096-4959(02)00073-8.
- The UniProt Consortium. 2017. UniProt: the universal protein knowledgebase. *Nucleic Acids Research*. 45:D158–D159 DOI:https://doi.org/10.1093/nar/gkw1099.
- Turk T, Kem WR. 2009. The phylum Cnidaria and investigations of its toxins and venoms until 1990. *Toxicon* 54:1031–1037 DOI:10.1016/j.toxicon.2009.06.031.
- Undheim EB, Jones A, Clauser KR, Holland JW, Pineda SS, King GF, Fry BG. 2014. Clawing through evolution: Toxin diversification and convergence in the ancient lineage Chilopoda (centipedes). *Molecular Biology and Evolution* 31:2124–2148 DOI: 10.1093/molbev/msu162.
- Uri S, Marina G, Liubov G. 2005. Severe delayed cutaneous reaction due to Mediterranean jellyfish (*Rhopilema nomadica*) envenomation. *Contact Dermatitis* 52:282–283 DOI: 10.1111/j.0105-1873.2005.00582.x.

- 630 Van Iten H, Marques AC, Leme JDM, Pacheco MLF, Simões MG. 2014. Origin and
631 early diversification of the phylum Cnidaria Verrill: Major developments in the analysis
632 of the taxon's Proterozoic-Cambrian history. *Palaeontology* 57:677–690 DOI:
633 10.1111/pala.12116.
- 634 Van Iten H, Leme JM, Pacheco MLAF, Simões MG, Fairchild TR, Rodrigues F, Galante
635 D, Boggiani PC, Marques AC. 2016. Origin and early diversification of phylum
636 Cnidaria: key macrofossils from the Ediacaran system of North and South America. In:
637 The Cnidaria, past, present and future, Goffredo, S. & Dubinsky, Z. (Ed.), Springer. pp.
638 31-40 (ISBN 978-3-319-31303-0; ISBN 978-3-319-31305-4 – e-book; 10.1007/978-3-
639 319-31305-4_3).
- 640 Weston AJ, Chung R, Dunlap WC, Morandini AC, Marques AC, Moura-da-Silva AM,
641 Ward M, Padilla G, da Silva LF, Andreakis N, Long PF. 2013. Proteomic
642 characterisation of toxins isolated from nematocysts of the South Atlantic jellyfish
643 *Olindias sambaquiensis*. *Toxicon* 71:11–17 DOI:10.1016/j.toxicon.2013.05.002.
- 644 Weston AJ, Dunlap WC, Shick JM, Klueter A, Iglic K, Vukelic A, Starcevic A, Ward M,
645 Wells ML, Trick CG, Long PF. 2012. A profile of an endosymbiont-enriched fraction of
646 the coral *Stylophora pistillata* reveals proteins relevant to microbial-host interactions.
647 *Molecular and Cellular Proteomics* 11 DOI: 10.1074/mcp.M111.015487.
- 648 Winkel KD, Tibballs J, Molenaar P, Lambert G, Coles P, Ross-Smith M, Wiltshire C,
649 Fenner PJ, Gershwin LA, Hawdon GM, Wright CE, Angus JA. 2005. Cardiovascular
650 actions of the venom from the Irukandji (*Carukia barnesi*) jellyfish: Effects in human,
651 rat and guinea-pig tissues *in vitro* and in pigs *in vivo*. *Clinical and Experimental*
652 *Pharmacology and Physiology* 32:777–788 DOI: 10.1111/j.1440-1681.2005.04258.x.
- 653

Figure legends

Figure 1: A) *Chrysaora lactea*, B) *Tamoya haplonema* and C) *Chiropsalmus quadrumanus*. Medusa adult stages of Cnidaria from which the venom proteomes of isolated nematocysts were acquired for this study. (Photos courtesy of Dr Alvaro Migotto, Centro de Biologia Marinha, Universidade de São Paulo São Sebastião, Brasil).

Figure 2. Comparison of Cnidarian venom composition. Venn diagram showing the number of putative toxin protein families shared among the soluble nematocyst proteomes of the four classes of cnidarians studied (Note that the protein families marked with an asterisk are described here for the first time).

Figure 3. Recruitment patterns of putative toxin protein families into Cnidaria venom, based on a established cnidarian phylogenies (Marques & Collins, 2004; Collins *et al.*, 2006). Solid black rectangles represent recruitment events. Dotted rectangles represent absence of toxin families. White rectangles represent multiple recruitments of toxin families. The numbers above of the lines represents the toxin families: 1. actinoporins; 2. complement C3; 3. conotoxins O; 4. Conotoxins T; 5. CRISP; 6. ficolin lectin; 7. flavin monoamine oxidase; 8. Jellyfish toxin; 9. kunitz-type; 10. latrotoxin; 11. MAC-PF; 12. metalloproteinase; 13. natriuretic peptide; 14. peptidase S1; 15. phospholipase A2; 16. phospholipase B; 17. phospholipase D; 18. ShK-like potassium channel; 19. snaclec; 20. snake three finger; 21. Sea anemone sodium channel modulator; 22.TCTP; 23. glycosyl hydrolase 56*; 24. huwentoxin-1*; 25. laticin*; 26. lipase*; 27. CS $\alpha\beta$ potassium channel blocker*; 28. CS $\alpha\beta$ sodium channel inhibitor*. The proteins families marked with asterisk (*) have never previously been recorded in Cnidaria.

Table 1: Predicted venom proteomes of potential toxins isolated from nematocysts. A) *Chiropsalmus quadrumanus*, B) *Tamoya haplonema* and C) *Chrysaora lactea*. Peptide fragments used for putative toxin annotation are given with validated spectra in Figures S1-S3.

| Toxin with closest homology | Predicted toxin protein family | Uniprot accession number | Example of animal species with closest homology |
|------------------------------------|---|--------------------------|--|
| A) <i>Chiropsalmus quadrumanus</i> | | | |
| Alpha-latroinsectotoxin-Lt1a | Latrotoxin | Q02989 | <i>Latrodectus tredecimguttatus</i> (European black widow spider) |
| Conotoxin Bu2 | Conotoxin O1 | P0CY61 | <i>Conus bullatus</i> (Bubble cone snail) |
| Echotoxin-2 | Actinoporin | Q76CA2 | <i>Monoplex parthenopeus</i> (Giant triton sea snail) |
| Hainantoxin-XVIII-5 | Putative ion channel inhibitor | D2Y2N9 | <i>Haplopelma hainanum</i> (Chinese bird spider) |
| Neurotoxin LmNaTx1 | CS $\alpha\beta$ sodium channel inhibitor | D9U297 | <i>Lychas mucronatus</i> (Chinese swimming scorpion) |
| Toxin CfTX-2 | Jellyfish toxin | A7L036 | <i>Chironex fleckeri</i> (Sea wasp) |
| B) <i>Tamoya haplonema</i> | | | |
| Alpha-latroinsectotoxin-Lt1a | Latrotoxin | Q02989 | <i>Latrodectus tredecimguttatus</i> (European black widow spider) |
| Conotoxin Lt5.9 | Conotoxin T | Q1A3Q7 | <i>Conus litteratus</i> (Lettered cone snail) |
| DELTA-alicitoxin-Pse2b | MACPF | P58912 | <i>Phyllodiscus semoni</i> (Wasp sea anemone) |
| Disintegrin acostatin-alpha | Disintegrin | Q805F7 | <i>Agkistrodon contortrix contortrix</i> (northern copperhead pit viper) |
| Echotoxin-2 | Actinoporin | Q76CA2 | <i>Monoplex parthenopeus</i> (Giant triton sea snail) |
| Equinatoxin-3 | Actinoporin | P0C1H2 | <i>Actinia equina</i> (Beadlet sea anemone) |

| | | | |
|--|--|--------|---|
| Im-conomorphin | Conotoxin A | P0CH39 | <i>Conus imperialis</i> (Imperial cone snail) |
| Maximins 3/H2 | Bombinin | P83082 | <i>Bombina maxima</i> (Yunnan firebelly toad) |
| Phospholipase A2 3 | Phospholipase A2 | P21792 | <i>Micrurus nigrocinctus</i> (Central American coral snake) |
| Phospholipase D L1SicTox-alphaIII2 | Arthropod phospholipase D | Q1KY79 | <i>Loxosceles laeta</i> (Chilean recluse spider) |
| Potassium channel toxin alpha-KTx Tx773 | CS $\alpha\beta$ potassium channel blocker | B8XH45 | <i>Buthus occitanus israelis</i> (Common yellow scorpion) |
| Potassium channel toxin TdiKIK | Long chain scorpion toxin | Q0GY43 | <i>Tityus discrepans</i> (Venezuelan scorpion) |
| Snake venom metalloproteinase aculysin-1 | Venom metalloproteinase (M12B) | Q9W7S2 | <i>Deinagkistrodon acutus</i> (Sharp-nosed pit viper) |
| U5-ctenitoxin-Co1a | Spider toxin Tx2 | P85276 | <i>Ctenus ornatus</i> (Brazilian spider) |
| Venom allergen 5 | CRISP | A9QQ26 | <i>Lycosa singoriensis</i> (Chinese wolf spider) |
| Venom carboxylesterase-6 | Lipase | B2D0J5 | <i>Apis mellifera</i> (European honey bee) |
| Venom nerve growth factor 1 | NGF-beta | Q2XXL6 | <i>Azemiops feae</i> (Black-headed Burmese viper) |

C) *Chrysaora lactea*

| | | | |
|---------------------------------------|--------------------------|--------|---|
| CfTX-2 | Jellyfish toxin | A7L036 | <i>Chironex fleckeri</i> (Sea wasp) |
| Cathelicidin-related peptide Na_CRAMP | Cathelicidin | B6S2X0 | <i>Naja atra</i> (Chinese cobra) |
| Conotoxin Bu2 | Conotoxin O1 | P0CY61 | <i>Conus bullatus</i> (Bubble cone snail) |
| Cysteine-rich venom protein LIO1 | CRISP | Q2XXQ0 | <i>Erythrolamprus poecilogyrus</i> (Water snake) |
| L-amino-acid oxidase | Flavin monoamine oxidase | P0DI84 | <i>Vipera ammodytes</i> (Sand viper) |
| M-zodatoxin-Lt4a | Latacin | Q1ELU5 | <i>Lachesana tarabaevi</i> (Ant spider) |
| Snake venom serine protease KN2 | Peptidase S1 | Q71QJ0 | <i>Trimeresurus stejnegeri</i> (Chinese green tree viper) |


U16-lycotoxin-Ls1a
Venom peptide Ocy2

U16-lycotoxin
Not assigned


B6DD52
P86107


Lycosa singoriensis (Chinese wolf spider)
Opisthacanthus cayaporum (South American scorpion)











- Early diverging metazoans offer a phylogenetic anchor to study evolution of the venom trait.
- Venom proteomes of the scyphozoan *Chrysaora lactea* and two cubozoans *Tamoya haplonema* and *Chiropsalmus quadrumanus* are presented.
- Toxin recruitment and retention patterns do not always correlate with accepted phylogeny.
- Factors that drive toxin diversification independent of phylogeny merit closer scrutiny.